

1

FUNCTIONS

OVERVIEW Functions are fundamental to the study of calculus. In this chapter we review what functions are and how they are pictured as graphs, how they are combined and transformed, and ways they can be classified. We review the trigonometric functions, and we discuss misrepresentations that can occur when using calculators and computers to obtain a function's graph. The real number system, Cartesian coordinates, straight lines, parabolas, and circles are reviewed in the Appendices. We treat inverse, exponential, and logarithmic functions in Chapter 7.

1.1

Functions and Their Graphs

Functions are a tool for describing the real world in mathematical terms. A function can be represented by an equation, a graph, a numerical table, or a verbal description; we will use all four representations throughout this book. This section reviews these function ideas.

Functions; Domain and Range

The temperature at which water boils depends on the elevation above sea level (the boiling point drops as you ascend). The interest paid on a cash investment depends on the length of time the investment is held. The area of a circle depends on the radius of the circle. The distance an object travels at constant speed along a straight-line path depends on the elapsed time.

In each case, the value of one variable quantity, say y , depends on the value of another variable quantity, which we might call x . We say that " y is a function of x " and write this symbolically as

$$y = f(x) \quad (\text{"}y\text{ equals }f\text{ of }x\text{"}).$$

In this notation, the symbol f represents the function, the letter x is the **independent variable** representing the input value of f , and y is the **dependent variable** or output value of f at x .

DEFINITION A function f from a set D to a set Y is a rule that assigns a *unique* (single) element $f(x) \in Y$ to each element $x \in D$.

The set D of all possible input values is called the **domain** of the function. The set of all values of $f(x)$ as x varies throughout D is called the **range** of the function. The range may not include every element in the set Y . The domain and range of a function can be any sets of objects, but often in calculus they are sets of real numbers interpreted as points of a coordinate line. (In Chapters 13–16, we will encounter functions for which the elements of the sets are points in the coordinate plane or in space.)

Often a function is given by a formula that describes how to calculate the output value from the input variable. For instance, the equation $A = \pi r^2$ is a rule that calculates the area A of a circle from its radius r (so r , interpreted as a length, can only be positive in this formula). When we define a function $y = f(x)$ with a formula and the domain is not stated explicitly or restricted by context, the domain is assumed to be the largest set of real x -values for which the formula gives real y -values, the so-called **natural domain**. If we want to restrict the domain in some way, we must say so. The domain of $y = x^2$ is the entire set of real numbers. To restrict the domain of the function to, say, positive values of x , we would write " $y = x^2, x > 0$ ".

Changing the domain to which we apply a formula usually changes the range as well. The range of $y = x^2$ is $[0, \infty)$. The range of $y = x^2, x \geq 2$, is the set of all numbers obtained by squaring numbers greater than or equal to 2. In set notation (see Appendix 1), the range is $\{x^2 | x \geq 2\}$ or $\{y | y \geq 4\}$ or $[4, \infty)$.

When the range of a function is a set of real numbers, the function is said to be **real-valued**. The domains and ranges of many real-valued functions of a real variable are intervals or combinations of intervals. The intervals may be open, closed, or half open, and may be finite or infinite. The range of a function is not always easy to find.

A function f is like a machine that produces an output value $f(x)$ in its range whenever we feed it an input value x from its domain (Figure 1.1). The function keys on a calculator give an example of a function as a machine. For instance, the \sqrt{x} key on a calculator gives an output value (the square root) whenever you enter a nonnegative number x and press the \sqrt{x} key.

A function can also be pictured as an **arrow diagram** (Figure 1.2). Each arrow associates an element of the domain D with a unique or single element in the set Y . In Figure 1.2, the arrows indicate that $f(a)$ is associated with a , $f(x)$ is associated with x , and so on. Notice that a function can have the same *value* at two different input elements in the domain (as occurs with $f(a)$ in Figure 1.2), but each input element x is assigned a *single* output value $f(x)$.

EXAMPLE 1 Let's verify the natural domains and associated ranges of some simple functions. The domains in each case are the values of x for which the formula makes sense.



FIGURE 1.1 A diagram showing a function as a kind of machine.




FIGURE 1.2 A function from a set D to a set Y assigns a unique element of Y to each element in D .

Function	Domain (x)	Range (y)
$y = x^2$	$(-\infty, \infty)$	$[0, \infty)$
$y = 1/x$	$(-\infty, 0) \cup (0, \infty)$	$(-\infty, 0) \cup (0, \infty)$
$y = \sqrt{x}$	$[0, \infty)$	$[0, \infty)$
$y = \sqrt{4 - x}$	$(-\infty, 4]$	$[0, \infty)$
$y = \sqrt{1 - x^2}$	$[-1, 1]$	$[0, 1]$

Solution The formula $y = x^2$ gives a real y -value for any real number x , so the domain is $(-\infty, \infty)$. The range of $y = x^2$ is $[0, \infty)$ because the square of any real number is nonnegative and every nonnegative number y is the square of its own square root $y = (\sqrt{y})^2$ for $y \geq 0$.

The formula $y = 1/x$ gives a real y -value for every x except $x = 0$. For consistency in the rules of arithmetic, we cannot divide any number by zero. The range of $y = 1/x$, the set of reciprocals of all nonzero real numbers, is the set of all nonzero real numbers, since $y = 1/(1/y)$. That is, for $y \neq 0$ the number $x = 1/y$ is the input assigned to the output value y .

The formula $y = \sqrt{x}$ gives a real y -value only if $x \geq 0$. The range of $y = \sqrt{x}$ is $[0, \infty)$ because every nonnegative number is some number's square root (namely, it is the square root of its own square).

In $y = \sqrt{4 - x}$, the quantity $4 - x$ cannot be negative. That is, $4 - x \geq 0$, or $x \leq 4$. The formula gives real y -values for all $x \leq 4$. The range of $\sqrt{4 - x}$ is $[0, \infty)$, the set of all nonnegative numbers.

The formula $y = \sqrt{1 - x^2}$ gives a real y -value for every x in the closed interval from -1 to 1 . Outside this domain, $1 - x^2$ is negative and its square root is not a real number. The values of $1 - x^2$ vary from 0 to 1 on the given domain, and the square roots of these values do the same. The range of $\sqrt{1 - x^2}$ is $[0, 1]$. ■

Graphs of Functions

If f is a function with domain D , its **graph** consists of the points in the Cartesian plane whose coordinates are the input-output pairs for f . In set notation, the graph is

$$\{(x, f(x)) \mid x \in D\}.$$

The graph of the function $f(x) = x + 2$ is the set of points with coordinates (x, y) for which $y = x + 2$. Its graph is the straight line sketched in Figure 1.3.

The graph of a function f is a useful picture of its behavior. If (x, y) is a point on the graph, then $y = f(x)$ is the height of the graph above the point x . The height may be positive or negative, depending on the sign of $f(x)$ (Figure 1.4).




FIGURE 1.3 The graph of $f(x) = x + 2$ is the set of points (x, y) for which y has the value $x + 2$.




FIGURE 1.4 If (x, y) lies on the graph of f , then the value $y = f(x)$ is the height of the graph above the point x (or below x if $f(x)$ is negative).

x	$y = x^2$
-2	4
-1	1
0	0
1	1
$\frac{3}{2}$	$\frac{9}{4}$
2	4






FIGURE 1.5 Graph of the function in Example 2.

EXAMPLE 2 Graph the function $y = x^2$ over the interval $[-2, 2]$.

Solution Make a table of xy -pairs that satisfy the equation $y = x^2$. Plot the points (x, y) whose coordinates appear in the table, and draw a *smooth* curve (labeled with its equation) through the plotted points (see Figure 1.5). ■

How do we know that the graph of $y = x^2$ doesn't look like one of these curves?



To find out, we could plot more points. But how would we then connect *them*? The basic question still remains: How do we know for sure what the graph looks like between the points we plot? Calculus answers this question, as we will see in Chapter 4. Meanwhile we will have to settle for plotting points and connecting them as best we can.

Representing a Function Numerically

We have seen how a function may be represented algebraically by a formula (the area function) and visually by a graph (Example 2). Another way to represent a function is **numerically**, through a table of values. Numerical representations are often used by engineers and scientists. From an appropriate table of values, a graph of the function can be obtained using the method illustrated in Example 2, possibly with the aid of a computer. The graph consisting of only the points in the table is called a **scatterplot**.

EXAMPLE 3 Musical notes are pressure waves in the air. The data in Table 1.1 give recorded pressure displacement versus time in seconds of a musical note produced by a tuning fork. The table provides a representation of the pressure function over time. If we first make a scatterplot and then connect approximately the data points (t, p) from the table, we obtain the graph shown in Figure 1.6.

TABLE 1.1 Tuning fork data

Time	Pressure	Time	Pressure
0.00091	-0.080	0.00362	0.217
0.00108	0.200	0.00379	0.480
0.00125	0.480	0.00398	0.681
0.00144	0.693	0.00416	0.810
0.00162	0.816	0.00435	0.827
0.00180	0.844	0.00453	0.749
0.00198	0.771	0.00471	0.581
0.00216	0.603	0.00489	0.346
0.00234	0.368	0.00507	0.077
0.00253	0.099	0.00525	-0.164
0.00271	-0.141	0.00543	-0.320
0.00289	-0.309	0.00562	-0.354
0.00307	-0.348	0.00579	-0.248
0.00325	-0.248	0.00598	-0.035
0.00344	-0.041		




FIGURE 1.6 A smooth curve through the plotted points gives a graph of the pressure function represented by Table 1.1 (Example 3).

The Vertical Line Test for a Function

Not every curve in the coordinate plane can be the graph of a function. A function f can have only one value $f(x)$ for each x in its domain, so no vertical line can intersect the graph of a function more than once. If a is in the domain of the function f , then the vertical line $x = a$ will intersect the graph of f at the single point $(a, f(a))$.

A circle cannot be the graph of a function since some vertical lines intersect the circle twice. The circle in Figure 1.7a, however, does contain the graphs of *two* functions of x : the upper semicircle defined by the function $f(x) = \sqrt{1 - x^2}$ and the lower semicircle defined by the function $g(x) = -\sqrt{1 - x^2}$ (Figures 1.7b and 1.7c).

EXAMPLE 6 The function whose value at any number x is the smallest integer greater than or equal to x is called the least integer function or the ceiling function. It is denoted $\lceil x \rceil$. Figure 1.11 shows the graph. For positive values of x , this function might represent, for example, the cost of parking x hours in a parking lot which charges \$1 for each hour or part of an hour.


$$\begin{aligned} \lceil 2 \rceil &= 2, & \lceil 0.2 \rceil &= 0, & \lceil -0.3 \rceil &= -1 & \lceil -2 \rceil &= -2, \\ \lceil 2.4 \rceil &= 2, & \lceil 1.9 \rceil &= 1, & \lceil 0 \rceil &= 0, & \lceil -1.2 \rceil &= -2, \end{aligned}$$

EXAMPLE 5 The function whose value at any number x is the greatest integer less than or equal to x is called the greatest integer function or the integer floor function. It is denoted $\lfloor x \rfloor$. Figure 1.10 shows the graph. Observe that

the position of x is determined by $y = \lfloor x \rfloor$. The values of f are given by $y = -x$ when $0 \leq x \leq 1$, and $y = 1$ when $x > 1$. The function, however, is just one function whose domain is the entire set of real numbers (Figure 1.9).


$$f(x) = \begin{cases} 1, & x < 1, \\ x^2, & 0 \leq x \leq 1 \\ -x, & x > 0 \end{cases}$$

EXAMPLE 4 The function



(Example 4).

To graph the function $y = f(x)$ shown here, we apply different formulas to different parts of its domain. The graph consists of two parts of its domain: $y = x^2$ when $x < 0$ and $y = 1$ when $x \geq 0$.



and range $[0, \infty)$.

The absolute value function has domain $(-\infty, \infty)$



whose graph is given in Figure 1.8. The right-hand side of the equation means that the function equals x if $x \geq 0$, and equals $-x$ if $x < 0$. Here are some other examples.

$$|x| = \begin{cases} -x, & x > 0, \\ x, & x \leq 0 \end{cases}$$

Sometimes a function is described by using different formulas on different parts of its domain. One example is the absolute value function

Piecewise-Defined Functions

FIGURE 1.7 (a) The circle is not the graph of a function; it fails the vertical line test. (b) The upper semicircle is the graph of a function $f(x) = \sqrt{1 - x^2}$. (c) The lower semicircle is the graph of a function $g(x) = -\sqrt{1 - x^2}$.



(Example 5).

it provides an integer floor for x lies on or below the line $y = x$, so greatest integer function $y = \lfloor x \rfloor$ for x less than or equal to x is $y = x$, so

FIGURE 1.10 The graph of the greatest integer function $y = \lfloor x \rfloor$.




FIGURE 1.11 The graph of the least integer function $y = \lceil x \rceil$ lies on or above the line $y = x$, so it provides an integer ceiling for x (Example 6).

Increasing and Decreasing Functions

If the graph of a function *climbs* or *rises* as you move from left to right, we say that the function is *increasing*. If the graph *descends* or *falls* as you move from left to right, the function is *decreasing*.

DEFINITIONS Let f be a function defined on an interval I and let x_1 and x_2 be any two points in I .

1. If $f(x_2) > f(x_1)$ whenever $x_1 < x_2$, then f is said to be **increasing** on I .
2. If $f(x_2) < f(x_1)$ whenever $x_1 < x_2$, then f is said to be **decreasing** on I .

It is important to realize that the definitions of increasing and decreasing functions must be satisfied for *every* pair of points x_1 and x_2 in I with $x_1 < x_2$. Because we use the inequality $<$ to compare the function values, instead of \leq , it is sometimes said that f is *strictly increasing* or *decreasing* on I . The interval I may be finite (also called bounded) or infinite (unbounded) and by definition never consists of a single point (Appendix 1).

EXAMPLE 7 The function graphed in Figure 1.9 is decreasing on $(-\infty, 0]$ and increasing on $[0, 1]$. The function is neither increasing nor decreasing on the interval $[1, \infty)$ because of the strict inequalities used to compare the function values in the definitions. ■

Even Functions and Odd Functions: Symmetry

The graphs of *even* and *odd* functions have characteristic symmetry properties.

DEFINITIONS A function $y = f(x)$ is an

- even function of x** if $f(-x) = f(x)$,
odd function of x if $f(-x) = -f(x)$,

for every x in the function's domain.

The names *even* and *odd* come from powers of x . If y is an even power of x , as in $y = x^2$ or $y = x^4$, it is an even function of x because $(-x)^2 = x^2$ and $(-x)^4 = x^4$. If y is an odd power of x , as in $y = x$ or $y = x^3$, it is an odd function of x because $(-x)^1 = -x$ and $(-x)^3 = -x^3$.

The graph of an even function is **symmetric about the y -axis**. Since $f(-x) = f(x)$, a point (x, y) lies on the graph if and only if the point $(-x, y)$ lies on the graph (Figure 1.12a). A reflection across the y -axis leaves the graph unchanged.

The graph of an odd function is **symmetric about the origin**. Since $f(-x) = -f(x)$, a point (x, y) lies on the graph if and only if the point $(-x, -y)$ lies on the graph (Figure 1.12b). Equivalently, a graph is symmetric about the origin if a rotation of 180° about the origin leaves the graph unchanged. Notice that the definitions imply that both x and $-x$ must be in the domain of f .

EXAMPLE 8

$$f(x) = x^2 \quad \text{Even function: } (-x)^2 = x^2 \text{ for all } x; \text{ symmetry about } y\text{-axis.}$$

$$f(x) = x^2 + 1 \quad \text{Even function: } (-x)^2 + 1 = x^2 + 1 \text{ for all } x; \text{ symmetry about } y\text{-axis}$$

(Figure 1.13a).

$$f(x) = x \quad \text{Odd function: } (-x) = -x \text{ for all } x; \text{ symmetry about the origin.}$$

$$f(x) = x + 1 \quad \text{Not odd: } f(-x) = -x + 1, \text{ but } -f(x) = -x - 1. \text{ The two are not equal.}$$

Not even: $(-x) + 1 \neq x + 1$ for all $x \neq 0$ (Figure 1.13b). ■





FIGURE 1.12 (a) The graph of $y = x^2$ (an even function) is symmetric about the y -axis. (b) The graph of $y = x^3$ (an odd function) is symmetric about the origin.

Power Functions A function $f(x) = x^a$, where a is a constant, is called a **power function**. There are several important cases to consider:
 If the variable y is proportional to the reciprocal $1/x$, then sometimes it is said that y is **inversely proportional** to x (because $1/x$ is the multiplicative inverse of x).

DEFINITION Two variables y and x are **proportional** (to one another) if one is always a constant multiple of the other; that is, if $y = kx$ for some nonzero constant k .

FIGURE 1.14 (a) Lines through the origin with slope m . (b) A constant function with slope $m = 0$.




Linear Functions A function of the form $f(x) = mx + b$, for constants m and b , is called a **linear function**. Figure 1.14a shows an array of lines $f(x) = mx$ where $b = 0$, so these lines pass through the origin. The function $f(x) = x$ where $m = 1$ and $b = 0$ is called the **identity function**. Constant functions result when the slope $m = 0$ (Figure 1.14b). A linear function with positive slope whose graph passes through the origin is called a **proportionality relationship**.

A variety of important types of functions are frequently encountered in calculus. We identify and briefly describe them here.

Common Functions

FIGURE 1.13 (a) When we add the constant term 1 to the function $y = x^2$, the resulting function $y = x^2 + 1$ is still even and its graph is still symmetric about the y -axis. (b) When we add the constant term 1 to the function $y = x$, the resulting function $y = x + 1$ is no longer odd. The symmetry about the origin is lost (Example 8).



(a) $a = n$, a positive integer.

The graphs of $f(x) = x^n$, for $n = 1, 2, 3, 4, 5$, are displayed in Figure 1.15. These functions are defined for all real values of x . Notice that as the power n gets larger, the curves tend to flatten toward the x -axis on the interval $(-1, 1)$, and also rise more steeply for $|x| > 1$. Each curve passes through the point $(1, 1)$ and through the origin. The graphs of functions with even powers are symmetric about the y -axis; those with odd powers are symmetric about the origin. The even-powered functions are decreasing on the interval $(-\infty, 0]$ and increasing on $[0, \infty)$; the odd-powered functions are increasing over the entire real line $(-\infty, \infty)$.




FIGURE 1.15 Graphs of $f(x) = x^n$, $n = 1, 2, 3, 4, 5$, defined for $-\infty < x < \infty$.

(b) $a = -1$ or $a = -2$.

The graphs of the functions $f(x) = x^{-1} = 1/x$ and $g(x) = x^{-2} = 1/x^2$ are shown in Figure 1.16. Both functions are defined for all $x \neq 0$ (you can never divide by zero). The graph of $y = 1/x$ is the hyperbola $xy = 1$, which approaches the coordinate axes far from the origin. The graph of $y = 1/x^2$ also approaches the coordinate axes. The graph of the function f is symmetric about the origin; f is decreasing on the intervals $(-\infty, 0)$ and $(0, \infty)$. The graph of the function g is symmetric about the y -axis; g is increasing on $(-\infty, 0)$ and decreasing on $(0, \infty)$.




FIGURE 1.16 Graphs of the power functions $f(x) = x^a$ for part (a) $a = -1$ and for part (b) $a = -2$.

(c) $a = \frac{1}{2}, \frac{1}{3}, \frac{3}{2}$, and $\frac{2}{3}$.


The functions $f(x) = x^{1/2} = \sqrt{x}$ and $g(x) = x^{1/3} = \sqrt[3]{x}$ are the **square root** and **cube root** functions, respectively. The domain of the square root function is $[0, \infty)$, but the cube root function is defined for all real x . Their graphs are displayed in Figure 1.17 along with the graphs of $y = x^{3/2}$ and $y = x^{2/3}$. (Recall that $x^{3/2} = (x^{1/2})^3$ and $x^{2/3} = (x^{1/3})^2$.)

Polynomials A function p is a **polynomial** if

$$p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$$


where n is a nonnegative integer and the numbers $a_0, a_1, a_2, \dots, a_n$ are real constants (called the **coefficients** of the polynomial). All polynomials have domain $(-\infty, \infty)$. If the

FIGURE 1.19 Graphs of three rational functions. The straight red lines are called asymptotes and are not part of the graph.




$g(x) \neq 0$. The graphs of several rational functions are shown in Figure 1.19. and g are polynomials. The domain of a rational function is the set of all x for which $f(x) = p(x)/g(x)$, where p

FIGURE 1.18 Graphs of three polynomial functions.



three polynomials. Techniques to graph polynomials are studied in Chapter 4. Polynomials $p(x) = ax^3 + bx^2 + cx + d$ of degree 3. Figure 1.18 shows the graphs of as $p(x) = ax^2 + bx + c$, are called quadratic functions. Likewise, cubic functions are functions with $m \neq 0$ are polynomials of degree 1. Polynomials of degree 2, usually written leading coefficient $a_n \neq 0$ and $n > 0$, then n is called the degree of the polynomial. Linear

FIGURE 1.17 Graphs of the power functions $f(x) = x^a$ for $a = \frac{1}{2}, \frac{3}{2}, \frac{2}{3}$, and $\frac{3}{2}$.



Algebraic Functions Any function constructed from polynomials using algebraic operations (addition, subtraction, multiplication, division, and taking roots) lies within the class of **algebraic functions**. All rational functions are algebraic, but also included are more complicated functions (such as those satisfying an equation like $y^3 - 9xy + x^3 = 0$, studied in Section 3.7). Figure 1.20 displays the graphs of three algebraic functions.




FIGURE 1.20 Graphs of three algebraic functions.

Trigonometric Functions The six basic trigonometric functions are reviewed in Section 1.3. The graphs of the sine and cosine functions are shown in Figure 1.21.




FIGURE 1.21 Graphs of the sine and cosine functions.

Exponential Functions Functions of the form $f(x) = a^x$, where the base $a > 0$ is a positive constant and $a \neq 1$, are called **exponential functions**. All exponential functions have domain $(-\infty, \infty)$ and range $(0, \infty)$, so an exponential function never assumes the value 0. We study exponential functions in Section 7.3. The graphs of some exponential functions are shown in Figure 1.22.





FIGURE 1.22 Graphs of exponential functions.

11. Express the edge length of a cube as a function of the cube's diagonal length.

10. Express the side length of a square as a function of the length d of the square's diagonal. Then express the surface area and volume of the cube as a function of the diagonal length.

9. Express the area and perimeter of an equilateral triangle as a function of the triangle's side length x .

7. a. Finding Formulas for Functions



In Exercises 7 and 8, which of the graphs are graphs of functions of x , and which are not? Give reasons for your answers.

$$5. f(t) = \frac{3-t}{t^2 - 16}$$

$$6. G(t) = \sqrt[3]{t} - t$$

$$4. g(x) = \sqrt{x^2 - 3x}$$

$$3. F(x) = \sqrt{5x + 10}$$

$$2. f(x) = 1 - \sqrt{x}$$

$$1. f(x) = 1 + x^2$$

Funtions

In Exercises 1–6, find the domain and range of each function.

Exercises 1.1

Transcendental Functions These are functions that are not algebraic. They include the trigonometric, inverse trigonometric, exponential, and logarithmic functions, and many other functions as well. A particular example of a transcendental function is a **catenary**. Its graph has the shape of a cable, like a telephone line or electric cable, strung from one support to another and hanging freely under its own weight (Figure 1.24). The function defining the graph is discussed in Section 7.7.

FIGURE 1.23 Graphs of four logarithmic functions.
The Latin word *catena* means "chain."






FIGURE 1.24 Graph of a catenary or hanging cable. (The Latin word *catena*



Logarithmic Functions These are the **inverse functions** of the exponential functions, and the calculus of these functions is studied in Chapter 7. Figure 1.23 shows the graphs of four logarithmic functions with various bases. In each case the domain is $(0, \infty)$ and the range is $(-\infty, \infty)$. Logarithmic functions are positive for $x > 0$ and approach negative infinity as $x \rightarrow 0^+$.

Exercises 1.1 Functions and Their Graphs



In Exercises 7 and 8, which of the graphs are graphs of functions of x , and which are not? Give reasons for your answers.

$$5. f(t) = \frac{3-t}{t^2 - 16}$$

$$6. G(t) = \sqrt[3]{t} - t$$

$$4. g(x) = \sqrt{x^2 - 3x}$$

$$3. F(x) = \sqrt{5x + 10}$$

$$2. f(x) = 1 - \sqrt{x}$$

$$1. f(x) = 1 + x^2$$

Funtions

In Exercises 1–6, find the domain and range of each function.

12. A point P in the first quadrant lies on the graph of the function $f(x) = \sqrt{x}$. Express the coordinates of P as functions of the slope of the line joining P to the origin.
13. Consider the point (x, y) lying on the graph of the line $2x + 4y = 5$. Let L be the distance from the point (x, y) to the origin $(0, 0)$. Write L as a function of x .
14. Consider the point (x, y) lying on the graph of $y = \sqrt{x - 3}$. Let L be the distance between the points (x, y) and $(4, 0)$. Write L as a function of y .

Functions and Graphs

Find the domain and graph the functions in Exercises 15–20.

15. $f(x) = 5 - 2x$ 16. $f(x) = 1 - 2x - x^2$
 17. $g(x) = \sqrt{|x|}$ 18. $g(x) = \sqrt{-x}$
 19. $F(t) = t/|t|$ 20. $G(t) = 1/|t|$

21. Find the domain of $y = \frac{x+3}{4-\sqrt{x^2-9}}$.

22. Find the range of $y = 2 + \frac{x^2}{x^2+4}$.

23. Graph the following equations and explain why they are not graphs of functions of x .

a. $|y| = x$ b. $y^2 = x^2$

24. Graph the following equations and explain why they are not graphs of functions of x .

a. $|x| + |y| = 1$ b. $|x + y| = 1$

Piecewise-Defined Functions

Graph the functions in Exercises 25–28.





25. $f(x) = \begin{cases} x, & 0 \leq x \leq 1 \\ 2 - x, & 1 < x \leq 2 \end{cases}$

26. $g(x) = \begin{cases} 1 - x, & 0 \leq x \leq 1 \\ 2 - x, & 1 < x \leq 2 \end{cases}$

27. $F(x) = \begin{cases} 4 - x^2, & x \leq 1 \\ x^2 + 2x, & x > 1 \end{cases}$

28. $G(x) = \begin{cases} 1/x, & x < 0 \\ x, & 0 \leq x \end{cases}$

Find a formula for each function graphed in Exercises 29–32.



The Greatest and Least Integer Functions

33. For what values of x is

a. $\lfloor x \rfloor = 0$? b. $\lceil x \rceil = 0$?

34. What real numbers x satisfy the equation $\lfloor x \rfloor = \lceil x \rceil$?

35. Does $\lceil -x \rceil = -\lfloor x \rfloor$ for all real x ? Give reasons for your answer.

36. Graph the function

$$f(x) = \begin{cases} \lfloor x \rfloor, & x \geq 0 \\ \lceil x \rceil, & x < 0. \end{cases}$$

Why is $f(x)$ called the *integer part* of x ?

Increasing and Decreasing Functions

Graph the functions in Exercises 37–46. What symmetries, if any, do the graphs have? Specify the intervals over which the function is increasing and the intervals where it is decreasing.

37. $y = -x^3$ 38. $y = -\frac{1}{x^2}$

39. $y = -\frac{1}{x}$ 40. $y = \frac{1}{|x|}$

41. $y = \sqrt{|x|}$ 42. $y = \sqrt{-x}$

43. $y = x^3/8$ 44. $y = -4\sqrt{x}$

45. $y = -x^{3/2}$ 46. $y = (-x)^{2/3}$

Even and Odd Functions

In Exercises 47–58, say whether the function is even, odd, or neither. Give reasons for your answer.

47. $f(x) = 3$

48. $f(x) = x^{-5}$

49. $f(x) = x^2 + 1$

50. $f(x) = x^2 + x$

51. $g(x) = x^3 + x$

52. $g(x) = x^4 + 3x^2 - 1$

53. $g(x) = \frac{1}{x^2 - 1}$

54. $g(x) = \frac{x}{x^2 - 1}$

55. $h(t) = \frac{1}{t-1}$

56. $h(t) = |t^3|$


57. $h(t) = 2t + 1$

58. $h(t) = 2|t| + 1$

Theory and Examples

59. The variable s is proportional to t , and $s = 25$ when $t = 75$. Determine t when $s = 60$.


60. Kinetic energy. The kinetic energy K of a mass is proportional to the square of its velocity v . If $K = 12,960$ joules when $v = 18 \text{ m/sec}$, what is K when $v = 10 \text{ m/sec}$?
61. The variables r and s are inversely proportional, and $r = 6$ when $s = 4$. Determine s when $r = 10$.
62. Boyle's Law. Boyle's Law says that the volume V of a gas at constant temperature T and pressure P is given by $V = k/T$, where k is a constant. If $V = 1000 \text{ in}^3$ when $P = 23.4 \text{ lbs/in}^2$ and $V = 147 \text{ in}^3$ when $P = 14.7 \text{ lbs/in}^2$, find k .
63. A box with an open top is to be constructed from a rectangular piece of cardboard with dimensions 14×22 in. by cutting out equal squares of side x at each corner and then folding up the sides to form a rectangular box. Express the volume V of the box as a function of x .



65. a. $y = x^4$ b. $y = x^7$ c. $y = x^{10}$

In Exercises 65 and 66, match each equation with its graph. Do not use a graphing device, and give reasons for your answer.


72. Industrial costs. A power plant sits next to a river where the river is 800 ft wide. To lay a new cable from the plant to a point Q on the opposite side goes from the plant to a point P on the opposite side of the river and then across the river and $\$100$ per foot across the river and $\$180$ per foot along the land.



71. A pen in the shape of an isosceles right triangle tapers with legs of length x and hypotenuse of length h if it is to be built. If fencing costs $\$5/\text{ft}$ for the legs and $\$10/\text{ft}$ for the hypotenuse, write the total cost of construction as a function of h .

70. Three hundred books sell for $\$40$ each, resulting in a revenue of $(300)(\$40) = \$12,000$. For each $\$5$ increase in the price, 25 fewer books are sold. Write the revenue R as a function of the number x of $\$5$ increases.
69. For a curve to be symmetric about the x -axis, the point (x, y) must together to identify the values of x for which lie on the curve if and only if the point $(x, -y)$ lies on the curve. Explain why a curve that is symmetric about the x -axis is not the graph of a function, unless the function is $y = 0$.


66. a. $y = 5x$ b. $y = 5x$ c. $y = x^5$



- b. Express the area of the rectangle in terms of x .

- a. Express the y -coordinate of P in terms of x . (You might start by writing an equation for the line AB .)


64. The accompanying figure shows a rectangle inscribed in an isosceles right triangle whose hypotenuse is 2 units long.



63. A box with an open top is to be constructed from a rectangular piece of cardboard with dimensions 14×22 in. by cutting out equal squares of side x at each corner and then folding up the sides to form a rectangular box. Express the volume V of the box as a function of x .

62. Boyle's Law. Boyle's Law says that the volume V of a gas at constant temperature T and pressure P are inversely proportional. If $P = 14.7 \text{ lbs/in}^2$ when $V = 1000 \text{ in}^3$, then what is V when $P = 23.4 \text{ lbs/in}^2$ when V and P are inversely proportional. If $P = 14.7 \text{ lbs/in}^2$ when $V = 1000 \text{ in}^3$, then what is P when $V = 147 \text{ in}^3$?

61. The variables r and s are inversely proportional, and $r = 6$ when $s = 4$. Determine s when $r = 10$.
60. Kinetic energy. The kinetic energy K of a mass is proportional to the square of its velocity v . If $K = 12,960$ joules when $v = 18 \text{ m/sec}$, what is K when $v = 10 \text{ m/sec}$?



59. a. $y = 5x$ b. $y = 5x$ c. $y = x^5$

1.2

Combining Functions; Shifting and Scaling Graphs

In this section we look at the main ways functions are combined or transformed to form new functions.

Sums, Differences, Products, and Quotients

Like numbers, functions can be added, subtracted, multiplied, and divided (except where the denominator is zero) to produce new functions. If f and g are functions, then for every x that belongs to the domains of both f and g (that is, for $x \in D(f) \cap D(g)$), we define functions $f + g$, $f - g$, and fg by the formulas

$$\begin{aligned}(f+g)(x) &= f(x) + g(x). \\ (f-g)(x) &= f(x) - g(x). \\ (fg)(x) &= f(x)g(x).\end{aligned}$$

Notice that the $+$ sign on the left-hand side of the first equation represents the operation of addition of *functions*, whereas the $+$ on the right-hand side of the equation means addition of the real numbers $f(x)$ and $g(x)$.

At any point of $D(f) \cap D(g)$ at which $g(x) \neq 0$, we can also define the function f/g by the formula

$$\left(\frac{f}{g}\right)(x) = \frac{f(x)}{g(x)} \quad (\text{where } g(x) \neq 0).$$

Functions can also be multiplied by constants: If c is a real number, then the function cf is defined for all x in the domain of f by

$$(cf)(x) = cf(x).$$

EXAMPLE 1 The functions defined by the formulas

$$f(x) = \sqrt{x} \quad \text{and} \quad g(x) = \sqrt{1-x}$$

have domains $D(f) = [0, \infty)$ and $D(g) = (-\infty, 1]$. The points common to these domains are the points

$$[0, \infty) \cap (-\infty, 1] = [0, 1].$$

The following table summarizes the formulas and domains for the various algebraic combinations of the two functions. We also write $f \cdot g$ for the product function fg .

Function	Formula	Domain
$f + g$	$(f+g)(x) = \sqrt{x} + \sqrt{1-x}$	$[0, 1] = D(f) \cap D(g)$
$f - g$	$(f-g)(x) = \sqrt{x} - \sqrt{1-x}$	$[0, 1]$
$g - f$	$(g-f)(x) = \sqrt{1-x} - \sqrt{x}$	$[0, 1]$
$f \cdot g$	$(f \cdot g)(x) = f(x)g(x) = \sqrt{x(1-x)}$	$[0, 1]$
f/g	$\frac{f}{g}(x) = \frac{f(x)}{g(x)} = \sqrt{\frac{x}{1-x}}$	$[0, 1) \quad (x = 1 \text{ excluded})$
g/f	$\frac{g}{f}(x) = \frac{g(x)}{f(x)} = \sqrt{\frac{1-x}{x}}$	$(0, 1] \quad (x = 0 \text{ excluded})$

The graph of the function $f + g$ is obtained from the graphs of f and g by adding the corresponding y -coordinates $f(x)$ and $g(x)$ at each point $x \in D(f) \cap D(g)$, as in Figure 1.25. The graphs of $f + g$ and $f \cdot g$ from Example 1 are shown in Figure 1.26.

The functions $f \circ g$ and $g \circ f$ are usually quite different.
 To evaluate the composite function $g \circ f$ (when defined), we find $f(x)$ first and then
 $g(f(x))$. The domain of $g \circ f$ is the set of numbers x in the domain of f such that $f(x)$ lies
 in the domain of g .

FIGURE 1.28 Arrow diagram for $f \circ g$.





FIGURE 1.27 Two functions can be composed at
 domain of the other. The composite is denoted by
 x whenever the value of one function at x lies in the
 $f \circ g$.



The definition implies that $f \circ g$ can be formed when the range of g lies in the
 domain of f . To find $(f \circ g)(x)$, first find $g(x)$ and second find $f(g(x))$. Figure 1.27 pic-
 tures $f \circ g$ as a machine diagram and Figure 1.28 shows the composite as an arrow di-
 agram.

The domain of $f \circ g$ consists of the numbers x in the domain of g for which $g(x)$
 lies in the domain of f .

$$(f \circ g)(x) = f(g(x))$$

DEFINITION If f and g are functions, the composite function $f \circ g$ (" f com-
 posed with g ") is defined by

Composition is another method for combining functions.

Composite Functions

function $f \circ g$ (Example 1).

FIGURE 1.26 The domain of the function $f + g$ is
 the intersection of the domains of f and g , the
 interval $[0, 1]$ on the x -axis where these domains
 overlap. This interval is also the domain of the
 function $f \cdot g$ (Example 1).





FIGURE 1.25 Graphical addition of two
 functions.



EXAMPLE 2 If $f(x) = \sqrt{x}$ and $g(x) = x + 1$, find

- (a) $(f \circ g)(x)$ (b) $(g \circ f)(x)$ (c) $(f \circ f)(x)$ (d) $(g \circ g)(x)$.

Solution

Composite

	Domain
(a) $(f \circ g)(x) = f(g(x)) = \sqrt{g(x)} = \sqrt{x+1}$	$[-1, \infty)$
(b) $(g \circ f)(x) = g(f(x)) = f(x) + 1 = \sqrt{x} + 1$	$[0, \infty)$
(c) $(f \circ f)(x) = f(f(x)) = \sqrt{f(x)} = \sqrt{\sqrt{x}} = x^{1/4}$	$[0, \infty)$
(d) $(g \circ g)(x) = g(g(x)) = g(x) + 1 = (x+1) + 1 = x+2$	$(-\infty, \infty)$

To see why the domain of $f \circ g$ is $[-1, \infty)$, notice that $g(x) = x + 1$ is defined for all real x but belongs to the domain of f only if $x + 1 \geq 0$, that is to say, when $x \geq -1$. ■

Notice that if $f(x) = x^2$ and $g(x) = \sqrt{x}$, then $(f \circ g)(x) = (\sqrt{x})^2 = x$. However, the domain of $f \circ g$ is $[0, \infty)$, not $(-\infty, \infty)$, since \sqrt{x} requires $x \geq 0$.

Shifting a Graph of a Function

A common way to obtain a new function from an existing one is by adding a constant to each output of the existing function, or to its input variable. The graph of the new function is the graph of the original function shifted vertically or horizontally, as follows.




FIGURE 1.29 To shift the graph of $f(x) = x^2$ up (or down), we add positive (or negative) constants to the formula for f (Examples 3a and b).

Shift Formulas

Vertical Shifts

$$y = f(x) + k \quad \begin{array}{l} \text{Shifts the graph of } f \text{ up } k \text{ units if } k > 0 \\ \text{Shifts it down } |k| \text{ units if } k < 0 \end{array}$$

Horizontal Shifts

$$y = f(x + h) \quad \begin{array}{l} \text{Shifts the graph of } f \text{ left } h \text{ units if } h > 0 \\ \text{Shifts it right } |h| \text{ units if } h < 0 \end{array}$$


EXAMPLE 3

- Adding 1 to the right-hand side of the formula $y = x^2$ to get $y = x^2 + 1$ shifts the graph up 1 unit (Figure 1.29).
- Adding -2 to the right-hand side of the formula $y = x^2$ to get $y = x^2 - 2$ shifts the graph down 2 units (Figure 1.29).
- Adding 3 to x in $y = x^2$ to get $y = (x + 3)^2$ shifts the graph 3 units to the left (Figure 1.30).
- Adding -2 to x in $y = |x|$, and then adding -1 to the result, gives $y = |x - 2| - 1$ and shifts the graph 2 units to the right and 1 unit down (Figure 1.31). ■

Scaling and Reflecting a Graph of a Function

To scale the graph of a function $y = f(x)$ is to stretch or compress it, vertically or horizontally. This is accomplished by multiplying the function f , or the independent variable x , by an appropriate constant c . Reflections across the coordinate axes are special cases where $c = -1$.

FIGURE 1.34 Reflections of the graph $y = \sqrt{x}$ across the coordinate axes (Example 4c).



(c) Reflection: The graph of $y = -\sqrt{x}$ is a reflection across the y-axis (Figure 1.34). $y = \sqrt{-x}$ is a reflection of $y = \sqrt{x}$ across the x-axis, and

stretching may correspond to a vertical stretching by a different scaling factor. Likewise, a horizontal stretching may correspond to a horizontal compression by a factor of 3 (Figure 1.33). Note that $y = \sqrt{3x} = \sqrt{x}/\sqrt{3}$ so a horizontal stretching by a factor of 3 $y = \sqrt{x}$ by a factor of 3, and $y = \sqrt{x}/3$ is a horizontal compression by a factor of 1/3 compresses the graph by a factor of 3 (Figure 1.32).

(a) Vertical: Multiplying the right-hand side of $y = \sqrt{x}$ by 3 to get $y = 3\sqrt{x}$ stretches the graph vertically by a factor of 3, whereas multiplying by 1/3 compresses the

EXAMPLE 4 Here we scale and reflect the graph of $y = \sqrt{x}$.

$y = f(-x)$ Reflects the graph of f across the y-axis.
 $y = -f(x)$ Reflects the graph of f across the x-axis.

For $c = -1$, the graph is reflected:

$y = f(x/c)$ Compresses the graph of f horizontally by a factor of c .
 $y = f(cx)$ Stretches the graph of f horizontally by a factor of c .


$y = \frac{1}{c} f(x)$ Compresses the graph of f vertically by a factor of c .
 $y = cf(x)$ Stretches the graph of f vertically by a factor of c .

For $c > 1$, the graph is scaled:

Vertical and Horizontal Scaling and Reflecting Formulas

down (Example 3d).

FIGURE 1.31 Shifting the graph of $y = |x|$ 2 units to the right and 1 unit



To shift the graph to the right, we add a positive constant to x (Example 3c). To shift the graph to the left, we add a negative constant to x .




FIGURE 1.33 Horizontally stretching and compressing the graph $y = \sqrt{x}$ by a factor of 3 (Example 4b).





FIGURE 1.32 Vertically stretching and compressing the graph $y = \sqrt{x}$ by a factor of 3 (Example 4a).



EXAMPLE 5 Given the function $f(x) = x^4 - 4x^3 + 10$ (Figure 1.35a), find formulas to

- compress the graph horizontally by a factor of 2 followed by a reflection across the y -axis (Figure 1.35b).
- compress the graph vertically by a factor of 2 followed by a reflection across the x -axis (Figure 1.35c).




FIGURE 1.35 (a) The original graph of f . (b) The horizontal compression of $y = f(x)$ in part (a) by a factor of 2, followed by a reflection across the y -axis. (c) The vertical compression of $y = f(x)$ in part (a) by a factor of 2, followed by a reflection across the x -axis (Example 5).

Solution

- (a) We multiply x by 2 to get the horizontal compression, and by -1 to give reflection across the y -axis. The formula is obtained by substituting $-2x$ for x in the right-hand side of the equation for f :

$$\begin{aligned}y &= f(-2x) = (-2x)^4 - 4(-2x)^3 + 10 \\&= 16x^4 + 32x^3 + 10.\end{aligned}$$

- (b) The formula is

$$y = -\frac{1}{2}f(x) = -\frac{1}{2}x^4 + 2x^3 - 5.$$

Ellipses

Although they are not the graphs of functions, circles can be stretched horizontally or vertically in the same way as the graphs of functions. The standard equation for a circle of radius r centered at the origin is

$$x^2 + y^2 = r^2.$$

Substituting cx for x in the standard equation for a circle (Figure 1.36a) gives

$$c^2x^2 + y^2 = r^2. \quad (1)$$




FIGURE 1.36 Horizontal stretching or compression of a circle produces graphs of ellipses.

Exercises 1.2

Equation (3) is the **standard equation of an ellipse** with center at (h, k) . The geometric definition and properties of ellipses are reviewed in Section 11.6.

$$(3) \quad \frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1.$$

Substituting $x = h$ for x , and $y = k$ for y , in Equation (2) results in


is vertical. The center of the ellipse given by Equation (2) is the origin (Figure 1.37). where $a = r/c$ and $b = r$. If $a > b$, the major axis is horizontal; if $a < b$, the major axis is vertical. The major axis is the line segment joining the points $(\pm r/c, 0)$. In both cases, the major axis is the longer line segment.

$$(2) \quad \frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

If we divide both sides of Equation (1) by r^2 , we obtain

the major axis is the line segment joining the points $(\pm r/c, 0)$. In both cases, ellipse are reversed in Figure 1.36c: the minor axis is the line segment joining the points $(0, \pm r)$, and the major axis is the line segment joining the points $(\pm r/c, 0)$. The axes of the ellipse are perpendicular to the major axis; the minor axis is the line segment joining the points $(0, \pm r)$, and the major axis is the line segment joining the points $(\pm r/c, 0)$. In Figure 1.36b, the line segment joining the points $(\pm r/c, 0)$ is called the **major axis** of the ellipse; the minor axis is the line segment joining the points $(0, \pm r)$. The graph of Equation (1) is an ellipse centered at the origin with horizontal major axis of length $2r/c$ and vertical minor axis of length $2r$.

FIGURE 1.37 Graph of the ellipse $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$, $a > b$, where the major axis is horizontal.



- Algebraic Combinations**
- In Exercises 1 and 2, find the domains and ranges of f , g , f/g , and f/g .
- Let $f(x) = x - 3$, $g(x) = \sqrt{x}$, $h(x) = x^3$, and $j(x) = 2x$. Express each of the functions in Exercises 11 and 12 as a composite of f , g , h , and j .
- vouling one or more of f , g , h , and j .
- Let $f(x) = x - 3$, $g(x) = \sqrt{x}$, $h(x) = x^3$, and $j(x) = 2x$. Express each of the functions in Exercises 11 and 12 as a composite of f , g , h , and j .
10. $f(x) = \frac{3-x}{x+2}$, $g(x) = \frac{x^2+1}{x^2}$, $h(x) = \sqrt[3]{2-x}$
11. $a. y = \sqrt{x-3}$
 b. $y = 2\sqrt{x-3}$
 c. $y = x^{1/4}$
 d. $y = 4x$
12. $a. y = 2x-3$
 b. $y = (2x-6)^3$
 c. $y = \sqrt{(x-3)^3}$
 d. $y = \sqrt[3]{x-6}$
13. Copy and complete the following table.
- | | | | |
|--------------------|-----------------|--------------|------------------|
| | $g(x)$ | $f(x)$ | $(f \circ g)(x)$ |
| a. $x - 7$ | \sqrt{x} | $?$ | $?$ |
| b. $x + 2$ | $3x$ | $\sqrt{x-5}$ | $?$ |
| c. $?$ | $\sqrt{x-5}$ | $3x$ | $?$ |
| d. $\frac{x}{x-1}$ | \sqrt{x} | $x-1$ | $?$ |
| e. $?$ | $1+\frac{1}{x}$ | x | $?$ |
| f. $\frac{1}{x}$ | $?$ | x | $?$ |
14. $f(x) = 1$, $g(x) = \sqrt{x-1}$
15. $f(x) = 2$, $g(x) = x^2 + 1$
16. $f(x) = \sqrt{x+1}$, $g(x) = \sqrt{x-1}$
17. $f(x) = \sqrt{x+1}$, $g(x) = \sqrt{x-1}$
18. $f(x) = x$, $g(x) = x^2 + 1$
19. $f(x) = \sqrt{x+1}$, $g(x) = \frac{x+4}{1}$, $h(x) = \frac{x}{1}$
20. $f(x) = x - 3$, $g(x) = \sqrt{x}$, $h(x) = x^3$, and $j(x) = 2x$. Express each of the functions in Exercises 11 and 12 as a composite of f , g , h , and j .
21. $a. y = 2\sqrt{x-3}$
 b. $y = x^3/2$
 c. $y = x^9$
 d. $y = 2\sqrt{x-3}$
22. $a. y = 2x-3$
 b. $y = (2x-6)^3$
 c. $y = \sqrt{(x-3)^3}$
 d. $y = 4x$
23. $f(x) = x^2 - 3$, $g(x) = \sqrt{x-3}$, and $j(x) = 2\sqrt{x-3}$
24. $f(x) = x - 3$, $g(x) = \sqrt{x}$, $h(x) = x^3$, and $j(x) = 2x$. Express each of the functions in Exercises 11 and 12 as a composite of f , g , h , and j .
25. If $f(x) = x + 5$ and $g(x) = x^2 - 3$, find the following.
- a. $f(g(0))$
 b. $g(f(0))$
 c. $f(g(x))$
 d. $g(f(x))$
 e. $f(f(-5))$
 f. $g(f(2))$
 g. $f(f(x))$
 h. $g(g(x))$
26. If $f(x) = x - 1$ and $g(x) = 1/(x+1)$, find the following.
- a. $f(g(1/2))$
 b. $g(f(1/2))$
 c. $f(g(x))$
 d. $g(f(x))$
 e. $f(f(x))$
 f. $g(g(x))$
 g. $f(f(x))$
 h. $g(g(2))$
27. In Exercises 7–10, write a formula for $f \circ g \circ h$.
28. $f(x) = 3x + 4$, $g(x) = 2x - 1$, $h(x) = x^2$
29. $f(x) = x + 1$, $g(x) = 3x$, $h(x) = 4 - x$
30. $f(x) = 3x + 4$, $g(x) = 2x - 1$, $h(x) = 4 - x$
31. **Composites of Functions**
32. If $f(x) = x - 1$ and $g(x) = 1/(x+1)$, find the following.
- a. $f(g(x))$
 b. $g(f(x))$
 c. $f(g(0))$
 d. $g(f(0))$
 e. $f(g(x))$
 f. $g(f(x))$
 g. $f(f(x))$
 h. $g(g(x))$
33. If $f(x) = x^2 - 3$, find the following.
- a. $f(g(x))$
 b. $g(f(x))$
 c. $f(g(0))$
 d. $g(f(0))$
 e. $f(g(x))$
 f. $g(f(x))$
 g. $f(f(x))$
 h. $g(g(x))$
34. If $f(x) = 1$, $g(x) = 1 + \sqrt{x}$
35. If $f(x) = x^2 + 1$
36. If $f(x) = 1 + \sqrt{x-1}$, $g(x) = \sqrt{x-1}$
37. If $f(x) = x$, $g(x) = \sqrt{x-1}$
38. If $f(x) = \sqrt{x+1}$, $g(x) = \sqrt{x-1}$
39. If $f(x) = x$, $g(x) = \sqrt{x+1}$
40. If $f(x) = 1$, $g(x) = \sqrt{x}$

14. Copy and complete the following table.

$g(x)$	$f(x)$	$(f \circ g)(x)$
a. $\frac{1}{x-1}$	$ x $?
b. ?	$\frac{x-1}{x}$	$\frac{x}{x+1}$
c. ?	\sqrt{x}	$ x $
d. \sqrt{x}	?	$ x $

15. Evaluate each expression using the given table of values

x	-2	-1	0	1	2
$f(x)$	1	0	-2	1	2
$g(x)$	2	1	0	-1	0

- a. $f(g(-1))$ b. $g(f(0))$ c. $f(f(-1))$
d. $g(g(2))$ e. $g(f(-2))$ f. $f(g(1))$

16. Evaluate each expression using the functions

$$f(x) = 2 - x, \quad g(x) = \begin{cases} -x, & -2 \leq x < 0 \\ x - 1, & 0 \leq x \leq 2. \end{cases}$$

- a. $f(g(0))$ b. $g(f(3))$ c. $g(g(-1))$
d. $f(f(2))$ e. $g(f(0))$ f. $f(g(1/2))$

In Exercises 17 and 18, (a) write formulas for $f \circ g$ and $g \circ f$ and find the (b) domain and (c) range of each.

17. $f(x) = \sqrt{x+1}$, $g(x) = \frac{1}{x}$


18. $f(x) = x^2$, $g(x) = 1 - \sqrt{x}$

19. Let $f(x) = \frac{x}{x-2}$. Find a function $y = g(x)$ so that $(f \circ g)(x) = x$.


20. Let $f(x) = 2x^3 - 4$. Find a function $y = g(x)$ so that $(f \circ g)(x) = x + 2$.

Shifting Graphs

21. The accompanying figure shows the graph of $y = -x^2$ shifted to two new positions. Write equations for the new graphs.




22. The accompanying figure shows the graph of $y = x^2$ shifted to two new positions. Write equations for the new graphs.



23. Match the equations listed in parts (a)–(d) to the graphs in the accompanying figure.

- a. $y = (x - 1)^2 - 4$ b. $y = (x - 2)^2 + 2$
c. $y = (x + 2)^2 + 2$ d. $y = (x + 3)^2 - 2$



24. The accompanying figure shows the graph of $y = -x^2$ shifted to four new positions. Write an equation for each new graph.

